



Europäisches  
Patentamt


European  
Patent Office

Office européen  
des brevets

PCT/IB 04 / 00002

12.01.04

#2



Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-  
gen stimmen mit der  
ursprünglich eingereichten  
Fassung der auf dem näch-  
sten Blatt bezeichneten  
europäischen Patentanmel-  
dung überein.

The attached documents  
are exact copies of the  
European patent application  
described on the following  
page, as originally filed.

Les documents fixés à  
cette attestation sont  
conformes à la version  
initialement déposée de  
la demande de brevet  
européen spécifiée à la  
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

03290082.1

Der Präsident des Europäischen Patentamts;  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

R C van Dijk

BEST AVAILABLE COPY

PRIORITY  
DOCUMENT

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

Anmeldung Nr:  
Application no.: 03290082.1  
Demande no:

Anmeldetag:  
Date of filing: 13.01.03  
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Orsymonde (Organisation de synthèse  
Mondiale)  
5, rue Charles Martigny  
94700 Maisons Alfort  
FRANCE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:  
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.  
If no title is shown please refer to the description.  
Si aucun titre n'est indiqué se référer à la description.)

Novel method for preparing methyl 2-diphenylmethylsulfinylacetate

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)  
revendiquée(s)  
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/  
Classification internationale des brevets:

C07C315/00

Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of  
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL  
PT SE SI SK TR LI

The present invention relates to a novel method for preparing methyl 2-diphenylmethylsulfinylacetate (MDMSA).

MDMSA is disclosed as an intermediate compound in the synthesis of modafinil also known as 2-[(diphenylmethyl)sulfinyl]acetamide. Modafinil which is a synthetic acetamide with wake-promoting activity, is useful in the treatment of narcolepsy, among other disorders.

The inventors have now discovered a novel route for synthesizing MDMSA which is applicable at the industrial scale.

Advantageously, the MDMSA can be obtained in two or three steps, each being characterized by high yields.


In an advantageous embodiment, these steps may be carried out in the same reactor and the same solvent, without isolating the intermediate compounds.

The aim of the present invention is to provide an economical and efficient method for preparing MDMSA.

These aims and others are achieved by the present invention which relates to a method for preparing methyl 2-diphenylmethylsulfinylacetate comprising the steps of:

- (i) conversion of benzhydrol into methyldiphenylmethylthioacetate ; and
- (ii) conversion of methyldiphenylmethylthioacetate into methyl-2-diphenylmethylsulfinylacetate by oxidation.

Scheme 1 illustrates in general the steps used in this method :



Scheme 1

The reactions of steps (i) and (ii) used in this method are carried out in an appropriate solvent which may be easily chosen by persons skilled in the art, it

being understood that an appropriate solvent denotes a solvent which is nonreactive towards the starting reagents, the intermediates or the products, at the reaction temperature considered, it being possible for the latter to vary from the solidification point of the solvent to the boiling point of the solvent.

5 A given reaction may be carried out in a solvent or in a mixture of several solvents, the solvent(s) being generally chosen according to the type of reaction considered and the subsequent treatment of the reaction medium.

In a preferred embodiment, the solvent is an aprotic solvent.

10 By way of illustration and without limitation of aprotic solvents which may be suitable for the method according to the invention, there may be mentioned in particular chlorinated solvents, aromatic solvents, hydrocarbon solvents and ethereal solvents.

Among the chlorinated solvents, chloroform, dichloromethane or chlorobenzene may be mentioned in particular.

15 Among the appropriate aromatic solvents, there may be mentioned; for example, benzene, toluene and chlorobenzene.

As examples of appropriate hydrocarbon solvents, cyclohexane, pentane and hexane may be mentioned.

20 Solvents such as diethyl ether, tetrahydrofuran and dioxane are useful as ethereal solvents.

Step (i) and (ii) may be conducted separately, notably in different solvents, where each intermediate is independently isolated.


In an advantageous embodiment, the reaction steps are conducted in the same reactor and without isolation of any intermediates.

#### 25 STEP i)

In a preferred embodiment, the conversion of benzhydrol into methyl diphenylmethylthioacetate comprises two steps of :

- 30 a1) conversion of benzhydrol into a benzhydryl carboxylate ; and  
b1) conversion of benzhydryl carboxylate into methyl diphenylmethyl thioacetate.

Scheme 2 illustrates in general steps a1) and b1) used in this method :



Scheme 2

*Step a1)*

In one preferred embodiment, step a1) comprises reacting benzhydrol and an acid anhydride in the presence of an inorganic acid and in an appropriate solvent, preferably aprotic.

In the context of the present invention, dichloromethane is particularly preferred for its advantageous properties of extraction, distillation at low temperature, its nonflammability, its chemical neutrality and its capacity to be easily recycled in the context of this method.

Generally, the acid anhydride and the aprotic solvent are loaded simultaneously into the reactor, at a temperature of about 20°C.

The acid anhydrides for the purposes of the present description denote either symmetric anhydrides of monocarboxylic acids, mixed anhydrides, or cyclic anhydrides of polycarboxylic acids.

By way of examples of acid anhydrides which may be suitable in step a1) according to the invention, there may be mentioned in particular acetic anhydride, butyric anhydride and propanoic anhydride, acetic anhydride being particularly preferred.

The preferred reaction conditions are those which provide for the use of an equimolar quantity of acid anhydride relative to the benzhydrol in the aprotic solvent.

However, it is preferable to carry out the procedure in the presence of a slight excess of acid anhydride. An excessively high excess may indeed induce the formation of by-products in the subsequent steps of the method while an excessively small quantity is capable of slowing the kinetics of the oxidation reaction carried out in step (ii). Thus, the molar ratio of acid anhydride preferably varies between 1 and 1.2, even better between 1 and 1.1, and is advantageously 1.05, which corresponds to the optimum molar ratio of acid anhydride which

makes it possible to obtain a clean synthesis.

Preferably, the procedure is carried out in the presence of a volume of solvent varying from 1.5 to 5 volumes, optimally in the presence of about 2 volumes. Under these conditions, the dilution brought about promotes the formation of the carboxylate.

According to an equally preferred variant of step a), the inorganic acid used is chosen from hydrochloric acid, hydrobromic acid, o-phosphoric acid and sulphuric acid, sulphuric acid being particularly preferred, in particular in the form of a 96% aqueous solution.

Preferably, the procedure is carried out in the presence of a quantity of inorganic acid ranging from 0.02 to 0.3 molar equivalents relative to the benzhydrol, even better from 0.05 to 0.15 equivalents.

As a safety precaution, the inorganic acid is generally introduced at a temperature of about 0°C so as to control the exothermicity of the reaction.

The benzhydrol is then loaded at a sufficient temperature to allow a sufficiently rapid reaction kinetics, but not too high so as to avoid the formation of by-products such as benzhydryl ether. It is particularly preferable to carry out the procedure at a temperature of between -5°C and +5°C, even better between -2°C and +2°C for an introduction time ranging from 45 minutes to 2 hours, preferably in about 1 hour. This introduction time indeed makes it possible to control the exothermicity of the reaction and to limit the formation of by-products.

The benzhydrol is generally maintained at this temperature for a sufficient contact time to obtain a complete reaction, but not too high in order to avoid the degradation of the benzhydryl carboxylate. The expression "complete reaction", for the purposes of the present invention, is understood to mean a reaction leading to the production of the derived product with a conversion rate greater than 99.2%, and preferably greater than 99.5%. In general, a complete reaction is obtained after a contact time of 2 hours.

The benzhydryl carboxylate obtained can be immediately used in the next step, without intermediate isolation.

Step a1) can also be realized by any other appropriate method.

As an example, step a1) can be realized by reacting benzhydrol with a carboxylic acid, for instance :

- 2-methyl-butiric acid in combination with :

- o  $\text{SOCl}_2$ , pyridine, in benzene ; or
- o  $\text{H}_2\text{SO}_4$  in dichloromethane ; or
- o  $\text{TsOH}$  in benzene

5 as disclosed in the reference : Fujita S. and al., Bull. Chem. Soc. Jpn 1972, 45 : 2571-2574 ;

- or acetic acid in combination with potassium iodide as described in Strazzoli P. et al., Recl. Trav. Chim., The Netherlands, 1991 ; 1:5-12.

10 By way of example, step a1) can also be realized by reacting a carboxylic acid salt, for instance acetic acid sodium salt as disclosed in Herzig S., Justus Liebigs, Ann. Chem. ; 1921 ; 422:330.

By way of example, step a1) can also be realized by reacting benzhydrol with a carboxylic acid chloride, as for example acetyl chloride in the presence of  
15 triethylamine as disclosed in Roizel B. and al., Chem. Commun., 2000, 16:1507-1508.

As an example, step a1) can also be realized by reacting benzhydrol with acetic acid ethyl ester in the presence of  $\text{Ti}(\text{OC}_2\text{H}_5)_4$  (Schnurrenberger P. and al., Helv. Chim. Acta, 1982, 65(4) : 1197-1201) ; or of iron (III) perchlorate (ITP)  
20 (Kumar B. and al., Indian J. Chem. Sect. B, 1993, 32(2) : 292-293) ; or of  $\text{Fe}(\text{ClO}_4)_3$ ,  $\text{SiO}_2$  (Parmar and al., Synth. Commun., 1999, 29(1) : 139-144).

#### Step b1)

Step b1) can be performed by any appropriate method.

25 In a preferred embodiment, step b1) comprises bringing the solution obtained in step a1) into contact with methyl thioglycolate.

The methyl thioglycolate is generally introduced in about 10 minutes at  $0^\circ\text{C}$  but with no temperature restriction (it can then increase to about  $9^\circ\text{C}$ ). The reaction medium is then heated to a sufficiently high temperature to drive the  
30 reaction kinetics, but not too high so as to avoid the formation of by-products. Generally, the procedure is carried out at a temperature of between  $15^\circ\text{C}$  and  $25^\circ\text{C}$ , preferably between  $18^\circ\text{C}$  and  $22^\circ\text{C}$ , and contact is maintained at this temperature for a sufficient time to obtain complete reaction, with few by-

products; generally for 2 to 3 hours, preferably for a period of 2 hours.


The methyl diphenylthioacetate can be used in step (ii) without intermediate isolation.

In another preferred embodiment, step i) comprises two steps of :

a2) conversion of benzhydrol into benzhydryl carboxylic acid ;

b2) conversion of benzhydryl carboxylic acid into methyl diphenyl-methylthioacetate.

This method is illustrated by scheme 3 :



Scheme 3

#### Step a2)

Step a2) may be realized according to any appropriate method and notably according to conditions disclosed in : Dahlbom O., Acta Chem. Scand., 1948, 2 : 856-858 ; Carceller E. et al., J. Med. Chem., 1993, 36 : 2984-2997 ; Lisac S. et al., J. Organomet. Chem. 1996, 507 : 215-220 ; Okarvi S. et al., J. Labelled Compd. Radiopharm, 1997, 39 : 853-874 ; Patent Thomae GmbH DE 2812542, 1979, Chem. Abstract 1980 ; 92 ; 198165 ; Iskander, Y. et al., J. Chem. Soc., 1961, 2397-2402.

In a particular embodiment, step a2) is realized by reacting benzhydrol with thioacetic acid, in the presence of an organic or inorganic acid.

Preferably, the solvent is a protic solvent, more preferably a carboxylic acid and notably acetic acid.

Inorganic or organic acid is preferably chosen among hydrochloric acid,  $\text{POCl}_3$ , trifluoroacetic acid, hydrobromic acid, o-phosphoric acid, sulphuric acid,  $\text{POCl}_3$  and trifluoroacetic acid being particularly preferred.



Preferably, the reaction is realized at room temperature.


**Step b2)**

Esterification reaction of step b2) may be realized by any methods known from the person skilled in the art.

In another particular embodiment, step (i) comprises two steps of :

- a3) conversion of the hydroxyl group of benzhydrol into a leaving group ;
- b3) conversion of the obtained product into methyldiphenyl-methylthioacetate.

This method is illustrated by scheme 4 :



Scheme 4

**Step a3)**

The leaving group in step a3) means any group that can be removed easily by a nucleophilic reactant. Leaving groups may be selected from the group consisting of halogenes, such as chloro- and bromo- radicals, or sulfonyl groups such as methanesulfonyl - or *p*-toluenesulfonyl- radicals.

Step a3) may be realized by any methods known from the person skilled in the art.

As an example, hydroxyl group of benzhydrol may be converted into chloro- or bromo- radical by reacting benzhydrol with thionyl chloride or thionyl bromide.


As an example, hydroxyl group of benzhydrol may be converted into methanesulfonate group or into *p*-toluenesulfonate group by reacting benzhydrol respectively with methanesulfonyl chloride or *p*-toluenesulfonyl chloride.

**Step b3)**

In a preferred embodiment, step b3) is realized according to conditions of step b1).

Step b3) may also be realized by any other appropriate method.

In another preferred embodiment, step i) comprises reacting benzhydrol with methylthioglycolate in the presence of a metallic catalyst in a solvent, as illustrated by scheme 5 :



Scheme 5

Preferably, the metallic catalyst is chosen from  $\text{ZnCl}_2$ ,  $\text{ZnBr}_2$ ,  $\text{ZnI}_2$ ,  $\text{ZnI}_2$  being particularly preferred.

Preferably, the solvent is chosen from aprotic solvents, more preferably from halogenated solvent and particularly chlorinated solvents such as dichloromethane, dichloroethane.

**STEP ii)**

Inventors have now settled oxidation conditions that allow interestingly to control the conversion of methylthiodiphenylmethylethanoate into methyl-2-diphenylmethanesulfinylethanoate, notably formation of by-products, yields and kinetics of the reaction.

Various parameters used in step (ii) capable of influencing the efficiency of the reaction may be optimized, such as the quantity of oxidizing agent introduced, the dilution of the reaction medium, the reaction temperature, the contact time and the acidity of the reaction medium.

Thus, in general, an excessively small quantity of oxidizing agent means an incomplete reaction. Conversely, an excessively large excess promotes the formation of the dioxidized by-product, sulfone.

By way of illustration of oxidizing agents which may be suitable for the invention, there may be mentioned in particular oxone, potassium permanganate, sodium percarbonate, peroxides such as hydrogen peroxide, tert-butyl hydroperoxide and m-chloroperoxybenzoic acid, hydrogen peroxide being particularly preferred.

In a preferred embodiment, step ii) is performed in dichloromethane.

According to a preferred embodiment, the oxidizing agent used in step (ii) is hydrogen peroxide, preferably in the form of a 35% aqueous solution. Indeed, a lower titre causes a higher dilution, which can decrease the kinetics of the reaction.

Although a stoichiometric quantity of oxidizing agent is sufficient, it is preferably to carry out the procedure in the presence of a slight excess, preferably in the presence of a molar ratio of between 1 and 1.1.

The acidity of the medium results from the operating conditions of step (i). The reaction temperature can influence the kinetics of the oxidation reaction. Thus, it is preferable that the temperature is between 28°C and 37°C, the temperature range for which the reaction kinetics is particularly increased taking into account the acidity of the medium.

Preferably, a reaction temperature of between 28°C and 32°C is preferred. This temperature indeed makes it possible to have optimum control of the method, in particular the stopping point beyond which superoxidation becomes non-negligible.

The contact time in order to obtain a clean and complete reaction may vary according to the scale of operation and also according to the quantity of inorganic acid, in particular of sulfuric acid, present in the reaction medium at step b).

Preferably, the reaction is considered as being "complete" in step (ii) when the ratio  $R_1 = \text{nonoxidized derivative} / (\text{monooxidized} + \text{dioxidized} + \text{nonoxidized derivative}) < 0.5\%$ .

The expression "clean" reaction is understood to mean, for the purposes of the present description, a reaction in which the ratio  $R_2 = \text{dioxidized derivative} / (\text{monooxidized} + \text{dioxidized} + \text{nonoxidized derivative}) < 0.5\%$ .

The contact time necessary in order to obtain a clean and complete reaction may be determined using conventional analytical techniques which make

it possible to monitor the progress of the reaction, such as HPLC (High-Performance Liquid Chromatography), IR (Infrared) or NMR (Nuclear Magnetic Resonance).

In general, the contact time necessary and sufficient to obtain a clean and complete reaction is less than 35 hours, preferably less than or equal to 33 hours, so as to avoid the formation of by-products such as in particular diphenylmethylthioacetic acid, and greater than 20 hours, preferably greater than or equal to 25 hours.

Of course, it is within the capability of persons skilled in the art to adjust the contact time necessary for obtaining a complete reaction at the scale of operation considered.

An increase in the quantity of inorganic acid in the reaction medium can nevertheless make it possible to significantly reduce the oxidation reaction time.

Without wishing to be limited to any theory, a hypothesis which makes it possible to explain this unexpected effect is that the inorganic acid plays a catalyst role in the mechanism of oxidation by the oxidizing agent. By way of example, in the case of sulphuric acid, it is assumed that a reaction intermediate of the  $\text{H}_2\text{SO}_5$  type is formed in the medium, transferring oxygen either directly to the oxidizable species, or indirectly by an accelerated formation of peracetic acid.

Thus, according to a preferred variant of step (ii) of the method according to the invention, an additional quantity of inorganic acid, preferably of 0.02 to 0.3 molar equivalents, and more preferably of 0.05 to 0.15 molar equivalents is added to the reaction medium in step b1); generally prior to the introduction of the oxidizing agent. An acceleration of the reaction kinetics is then observed.

Advantageously, the contact time required which is sufficient to obtain a complete and clean reaction in step (ii) is considerably reduced and is generally between 10 and 13 hours.

Advantageously, the introduction of the inorganic acid in two portions makes it possible to reduce the acidity of the reaction medium in steps a1) and b1) and therefore to limit the formation of by-products.

#### STEP (iii)

In a further embodiment, the method according to the invention comprises

an additional step of iii) recovering the methyl 2-diphenyl-methylsulfinylacetate obtained.

The isolation of the MDMSA formed in step (iii) can be carried out according to any conventional method known to persons skilled in the art.

Preferably, the MDMSA is isolated by extraction.

The organic phases are then combined and concentrated under reduced pressure, preferably at a temperature of 70°C.

According to a particular variant, the solvent is distilled to dryness.

The product may be purified according to any method known to persons skilled in the art such as recrystallization or chromatography.

According to a particular embodiment, step (iii) may comprise a step of direct crystallization of the MDMSA.

The expression "direct crystallization", for the purposes of the present description, is understood to mean a crystallization of the noncrystallized product caused by the addition of appropriate solvents, preferably chosen in particular from methanol, ethanol, ethyl acetate, isopropyl acetate and toluene, isopropyl acetate being particularly preferred.

In this context, the crystallization solvent is introduced after substantial removal of the aprotic solvent.

Advantageously, this direct crystallization makes it possible to purify the crude product immediately in the remainder of the method and thus to dispense with an isolation step and a more costly subsequent retreatment step.


In a particularly preferred embodiment, the method comprises the steps of:

- i) a1) converting benzhydrol into a benzhydryl carboxylate by reacting benzhydrol and an acid anhydride in the presence of an inorganic acid and in an appropriate aprotic solvent ;  
b1) converting the benzhydryl carboxylate into methyl diphenylmethylthioacetate by bringing the above solution into contact with methyl thioglycolate;
- ii) converting the diphenylmethylthioacetate into methyl 2-diphenylmethylsulfinylacetate by bringing the above solution into contact with an oxidizing agent ;

and optionally ;

iii) recovering the methyl 2-diphenylmethylsulfinylacetate obtained.

This embodiment is illustrated by scheme 6 :



Scheme 6

Advantageously, the MDMSA is obtained in three successive steps, each being characterized by high yields.

Furthermore, these three steps may be carried out in the same reactor and the same solvent, without isolating the intermediate compounds.

Advantageously, the formation of undesirable by-products is limited and controlled according to this method, which makes it possible to dispense with subsequent reprocessing steps.

Finally, according to another advantageous aspect, this method makes it possible to reduce the costs of manufacturing MDMSA, in particular because of its ease of use, and its high productivity and reproducibility.

The present invention also aims at the MDMSA obtained by the method of the invention and particularly by the method implementing :

- i) a1) converting benzhydrol into a benzhydryl carboxylate by reacting benzhydrol and an acid anhydride in the presence of an inorganic acid and in an appropriate aprotic solvent ;
- b1) converting the benzhydryl carboxylate into methyl

diphenylmethylthioacetate by bringing the above solution into contact with methyl thioglycolate;

- ii) converting the diphenylmethylthioacetate into methyl 2-diphenylmethanesulfinylacetate by bringing the above solution into contact with an oxidizing agent ;  
and optionally ;
- iii) recovering the methyl 2-diphenylmethanesulfinylacetate obtained.

The present invention is also directed to a method for preparing modafinil comprising the steps of :

- (i) conversion of benzhydrol into methyldiphenylmethylthioacetate ; and
- (ii) conversion of methyldiphenylmethylthioacetate into methyl-2-diphenylmethanesulfinylacetate by oxidation.

EXAMPLES

Example 1. – Synthesis of MDMSA on the laboratory scale (0.5 l)

5 a) Benzhydryl acetate

108.3 g (1.05 mol; 1.05 eq) of acetic anhydride are diluted in 370 ml of dichloromethane at 20°C. The solution obtained is cooled to  $0 \pm 2^\circ\text{C}$  before introducing 2.8 ml of a 96% sulphuric acid solution in a time of about 10 minutes. After stirring for about 10 minutes, 184.2 g (1 mol, 1 eq) of benzhydrol are  
10 introduced in portions at  $0^\circ\text{C} \pm 2^\circ\text{C}$  in  $60 \pm 15$  minutes. The reaction medium is maintained in contact for 2 hours at this temperature.

b) Methyl diphenylmethylthioacetate (MDMTA)

108.3 g (1.02 mol; 1.02 eq) of methyl thioglycolate are introduced at  $0^\circ\text{C} \pm$   
15  $2^\circ\text{C}$ , and then the reaction mixture is heated to  $20^\circ\text{C} \pm 2^\circ\text{C}$  and kept in contact for 2 hours at this temperature.

c) Methyl 2-diphenylmethylsulfinylacetate (MDMSA)

The reaction medium is heated to  $30^\circ\text{C} \pm 2^\circ\text{C}$ , and  $100.5 \pm 0.5$  g of a 35%  
20 hydrogen peroxide solution are added. The progress of the oxidation reaction is monitored by HPLC. After 25 hours of contact, the reaction is complete ( $R1$  and  $R2 < 0.5\%$ ).

d) Recovery of MDMSA

25 150 ml of distilled water are stirred. A 25% aqueous ammonia solution is added so as to bring the pH to 8. The aqueous organic phases are separated, and the aqueous phase is extracted with 2 x 100 ml of dichloromethane. The organic phases are then combined and concentrated under reduced pressure at a temperature of  $70^\circ\text{C}$ . The concentrate obtained crystallizes to give, after  
30 grinding, a white powder with a yield of  $98.0 \pm 0.5\%$ .



Examples 2 and 3. – Synthesis of MDMSA on a pilot scale (100 l)

*Example 2: Synthesis of crystallized MDMSA*

a) Benzhydryl acetate

5 A solution of acetic anhydride (8.73 kg; 85.5 mol; 1.05 eq) in methylene chloride (40 l) is prepared at  $20^{\circ}\text{C} \pm 2^{\circ}\text{C}$ . A 96% sulphuric acid solution (225 ml; 4.1 mol; 0.05 eq) is then added with a stirring of 100 revolutions/min, at  $0^{\circ}\text{C} \pm 5^{\circ}\text{C}$  in 5 to 10 minutes. The benzhydrol (15 kg; 81.4 mol, 1 eq) is then introduced at  $0^{\circ}\text{C} \pm 2^{\circ}\text{C}$  in 1.25 hours. The reaction mixture is stirred for a contact time of 2  
10 hours.

b) Methyl diphenylmethylthioacetate (MDMTA)

Methyl thioglycolate (8.81 kg; 82.9 mol; 1.02 eq) is introduced at  $0^{\circ}\text{C} \pm 2^{\circ}\text{C}$  in about 10 minutes. The reaction mixture is brought to the temperature of  $20^{\circ}\text{C} \pm 2^{\circ}\text{C}$  and kept at this temperature for a contact time of 2 hours, with a stirring of  
15 100 revolutions/min.

c) Methyl 2-diphenylmethylsulfinylacetate (MDMSA)

The reaction mixture is then brought to  $30^{\circ}\text{C}$  before introducing sulphuric  
20 acid (450 ml; 8.1 mol; 0.1 eq) with stirring (100 revolutions/min), in about 5 to 10 minutes. A 35% hydrogen peroxide solution (8.19 kg; 84.3 mol; 1.035 eq) is then introduced at  $30^{\circ}\text{C} \pm 2^{\circ}\text{C}$  in 1 hour. The contact time is determined by monitoring the reaction by HPLC (cf. Table 1).

25 d) Recovery of MDMSA

The mixture is cooled to  $20^{\circ}\text{C} \pm 2^{\circ}\text{C}$  and then 20 l of water are introduced. After neutralizing the reaction medium by adding a sufficient quantity of  $\text{NH}_4\text{OH}$  so that  $8 < \text{pH} < 9$ , the aqueous and organic phases are separated and the aqueous phase extracted twice with 10 l of methylene chloride. The chlorinated  
30 phases are washed with 10 l of water.

The solvent is distilled under atmospheric pressure and then under reduced pressure, at a jacket temperature of  $70^{\circ}\text{C}$ . When the distillation is

complete, isopropyl acetate (1.8 vol; 42 l) is added and the whole is cooled to -10°C. After draining and drying under vacuum at 45°C, MDMSA is obtained.

*Table 1: Yield and quality of the MDMSA and of the intermediate products obtained by this method for trials 1 to 4 (step (ii)).*

Trial	Formation of benzhydryl acetate		Formation of MDMA		Formation of MDMSA		HPLC of the finished product	Yield
	Time	HPLC %	Time	HPLC %	Time	HPLC %	%	
1	2h	98.9	2 h	98.5	10h	T = 1 S = 0.3	D = 99.7 S = 0.3	88
					11h	T = 0.7 S = 0.35		
					11h30	T = 0.3 S = 0.4		
2	2h	99.3	2h	99.1	11h30	T = 0.9 S = 0.35	D = 99.6 S = 0.3 T = 0.1	86.4
3	2h	99.3	2h	99.0	11h30	T = 0 S = 0.4	D = 99.7 S = 0.3	90.1
4	2h	99.3	2h	99.1	5 h	T = 7.2 S = 0.16	D = 99.7 S = 0.3	90.0
					6h30	T = 4.4 S = 0.17		
					8h	T = 2.3 S = 0.22		
					9h	T = 1.6 S = 0.14		
					10h	T = 1.15 S = 0.3		
					11h	T = 0.6 S = 0.38		
					11h30	T = 0.3 S = 0.38		

D = MDMSA

S = Sulphone

T = MDMA

10

These results demonstrate that the production of benzhydryl acetate and of MDMA can be reproducible.

The formation of MDMSA takes place in 11 h 30 min and reproducibly gives, after crystallization from isopropyl acetate, a final product which is in conformity ( $R_1$  and  $R_2 < 0.5\%$ ), with a yield of the order of 90%.

5      *Example 3: Synthesis of crude MDMSA*

a) Benzhydryl acetate

A solution of acetic anhydride (8.73 kg; 85.5 mol; 1.05 eq) in methylene chloride (40 l) is prepared at 20°C. A 96% sulphuric acid solution (225 ml; 4.1 mol; 0.05 eq) is then added with a stirring of 100 revolutions/min, at 0°C ± 5°C in 5 to 10 minutes. The benzhydrol (15 kg; 8.4 mol, 1 eq) is then introduced at 0°C ± 2°C in 1.25 hours. The reaction mixture is stirred for a contact time of 2 hours.

b) Methyl diphenylmethylthioacetate (MDMTA)

15      Methyl thioglycolate (8.81 kg; 82.9 mol; 1.02 eq) is introduced at 0°C in about 10 minutes. The reaction mixture is brought to the temperature of 20°C ± 2°C and kept at this temperature for a contact time of 2 hours, with stirring of 100 revolutions/min.

20      c) Methyl 2-diphenylmethylsulfinylacetate (MDMSA)

Once the reaction mixture has been brought to 30°C ± 2°C, a 35% hydrogen peroxide solution (8.19 kg; 84.3 mol; 1.035 eq) is introduced with stirring (100 revolutions/min) in 1 hour. The contact time is determined by monitoring the reaction by HPLC (cf. Table 2).

25      d) Recovery of MDMSA

The mixture is cooled to 20°C and then 20 l of water are introduced. After neutralizing the reaction medium by adding a sufficient quantity of  $\text{NH}_4\text{OH}$  so that  $8 < \text{pH} < 9$ , the aqueous and organic phases are separated and the aqueous phase extracted twice with 10 l of methylene chloride. The chlorinated phases are washed with 10 l of water.

30      The solvent is distilled to dryness under atmospheric pressure and then under reduced pressure at a jacket temperature of 70°C in a Moritz® turbosphere.

*Table 2: Yield and quality of the finished product and of the intermediate products for trials 5 and 6*

Trial	Formation of benzhydryl acetate		Formation of MDMA		Formation of MDMSA		HPLC of the finished product	Yield
	Time	HPLC %	Time	HPLC %	Time	HPLC %	%	%
5	2h	99.6	2h	99.2	35h	T = 0.3 S = 0.4	D = 99.12 S = 0.44 T = 0.24	97.2
6	2h	99.3	2h	98.9	33h	T = 0.17 S = 0.4	D = 99.2 S = 0.4 T = 0.2	97

5 D = MDMSA

S = Sulphone

T = MDMA

10 These results show that the steps of formation of benzhydryl acetate and of MDMA can be reproducible.

The step of oxidation of MDMA requires a contact time of about 33-35 hours and gives an MDMSA product which is in conformity ( $R_1$  and  $R_2 < 0.5\%$ ), with good yields (of the order of 97%).

CLAIMS

1. Method for preparing methyl 2-diphenylmethylsulfinylacetate (MDMSA) comprising the steps of :

- 5           (i) conversion of benzhydrol into methyldiphenylmethylthioacetate ; and
- (ii) conversion of methyldiphenylmethylthioacetate into methyl-2-diphenylmethylsulfinylacetate.

10           2. Method according to claim 1, in which step (i) comprises the following steps :

- a1) conversion of benzhydrol to benzhydrol carboxylate in an appropriate solvent ;
- b1) conversion of the benzhydrol carboxylate to methyl
- 15           diphenylmethylthioacetate.

          3. Method according to claim 2, in which the step (a1) comprises reacting benzhydrol and an acid anhydride in the presence of an inorganic acid and in an appropriate solvent.

20           4. Method according to claim 3, in which the solvent is an aprotic solvent.

          5. Method according to claim 4, in which the aprotic solvent is chosen from chlorinated solvents, aromatic solvents, hydrocarbon solvents and ethereal

25           solvents.

          6. Method according to claim 5, in which the aprotic solvent is chosen from chlorinated solvents.

30           7. Method according to claim 6, in which the solvent is dichloromethane.

8. Method according to any one of claims 3 to 7, in which the acid anhydride is chosen from acetic anhydride, propanoic anhydride and butyric anhydride.

5 9. Method according to claim 8, in which the acid anhydride is acetic anhydride.

10 10. Method according to any one of claims 3 to 9, in which the inorganic acid is chosen from hydrochloric acid, butyric acid, o-phosphoric acid and sulfuric acid.

11. Method according to claim 10, in which the inorganic acid is sulfuric acid.

15 12. Method according to any one of claims 3 to 11, in which the quantity of inorganic acid used is from 0.02 to 0.3 molar equivalents relative to the benzhydrol.

20 13. Method according to any one of claims 3 to 12, in which the reaction temperature in step a) is between  $-5^{\circ}\text{C}$  and  $+5^{\circ}\text{C}$ .

14. Method according to claim 2 to 13, in which step b1) comprises bringing the solution obtained in step a) into contact with methyl thioglycolate.

25 15. Method according to claim 14, in which the contact time used in step b1) is between 2 and 3 hours.

30 16. Method according to claim 14 to 15, in which the contact temperature used in step b1) is between  $15^{\circ}\text{C}$  and  $25^{\circ}\text{C}$ .

17. Method according to any one of the preceding claims, in which the oxidizing agent is chosen from oxone, potassium permanganate, sodium

percarbonate, peroxides such as hydrogen peroxide, tert-butyl hydroperoxide and m-chloroperoxybenzoic acid.

18. Method according to claim 17, in which the oxidizing agent is hydrogen  
5 peroxide.

19. Method according to claim 18, in which the hydrogen peroxide is added in the form of a 35% aqueous solution.

10 20. Method according to any one of the preceding claims, in which the oxidizing agent is used in an amount of 1 to 1.1 molar equivalent.

21. Method according to any one of the preceding claims, in which the reaction temperature in step (ii) is between 28°C and 37°C.  
15

22. Method according to one of claims 3 to 21, in which an additional quantity of inorganic acid is added in step (ii).

23. Method according to claim 22, in which the additional quantity of  
20 inorganic acid is from 0.02 to 0.3 molar equivalents.

24. Method according to either of claims 22 and 23, in which the contact time in step (ii) is between 10 and 13 hours.

25 25. Method according to any one of the preceding claims, which comprises an additional step (iii) recovering the methyl 2-diphenyl-methylsulfinylacetate obtained.

26. Method according to claim 25, in which step (iii) comprises a distillation  
30 of the solvent to dryness.

27. Method according to any one of claims 25 to 26, in which step (iii) comprises a step of direct crystallization.

28. Method according to claim 27, in which the crystallization solvent is chosen from methanol, ethanol, ethyl acetate, isopropyl acetate and toluene.

5 29. Method according to claim 28, in which the crystallization solvent is isopropyl acetate.

30. Method according to any one of the preceding claims, in which the successive steps are carried out in the same reactor without isolation of the  
10 intermediate compounds.

31. Method for preparing modafinil comprising preparing MDMSA according to claims 1 to 30.



ORSYMONDE

Novel method for preparing  
methyl 2-diphenylmethylsulfinylacetate

ABSTRACT

Method for preparing methyl 2-diphenylmethylsulfinylacetate (MDMSA) comprising the steps of :

- (i) conversion of benzhydrol into methyldiphenylmethylthioacetate ; and
- (ii) conversion of methyldiphenylmethylthioacetate into methyl-2-diphenylmethylsulfinylacetate by oxidation.

Figure: none.

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**